

Nonideal Reaction-Diffusion Systems: Multiple Routes to Instability

Timur Aslyamov^{1,*}, Francesco Avanzini^{1,2,†}, Étienne Fodor^{1,‡}, and Massimiliano Esposito^{1,§}
¹*Department of Physics and Materials Science, University of Luxembourg, L-1511 Luxembourg City, Luxembourg*
²*Department of Chemical Sciences, University of Padova, Via F. Marzolo, 1, I-35131 Padova, Italy*

 (Received 21 April 2023; accepted 30 August 2023; published 27 September 2023)

We develop a general classification of the nature of the instabilities yielding spatial organization in open nonideal reaction-diffusion systems, based on linear stability analysis. This encompasses dynamics where chemical species diffuse, interact with each other, and undergo chemical reactions driven out of equilibrium by external chemostats. We find analytically that these instabilities can be of two types: instabilities caused by intermolecular energetic interactions (E type), and instabilities caused by multimolecular out-of-equilibrium chemical reactions (R type). Furthermore, we identify a class of chemical reaction networks, containing unimolecular networks but also extending beyond them, that can only undergo E-type instabilities. We illustrate our analytical findings with numerical simulations on two reaction-diffusion models, each displaying one of the two types of instability and generating stable patterns.

DOI: [10.1103/PhysRevLett.131.138301](https://doi.org/10.1103/PhysRevLett.131.138301)

Introduction.—Reaction-diffusion (RD) systems play a crucial role in explaining the emergence of many spatial structures across scales, e.g., spiral form of galaxies [1], predator-prey distributions in ecological models [2], skin color patterns of animals [3], self-organization at the molecular scale [4], phase separation in electrochemical batteries [5]. The foundation of RD theory dates back to the seminal paper of A. M. Turing [6], where he proposed a spatial symmetry-breaking mechanism yielding stationary patterns.


Subsequent studies by the Brussels school of thermodynamics, led by I. Prigogine, showed the physicochemical and thermodynamical relevance of Turing patterns. Since they considered ideal solutions where the concentration dynamics is governed by linear diffusion and mass-action kinetics, they emphasized the need to consider multimolecular reactions and open systems driven far from equilibrium to generate patterns [7–9]. Indeed, in ideal solutions, on the one hand, multimolecular reactions are necessary to generate purely entropic interactions between species which create the nonlinearities at the basis of the instabilities, and on the other hand, nonequilibrium drives are required to prevent relaxation toward homogeneous concentration profiles. Since then, RD structures in ideal solutions have been extensively studied [10–13].

Nonideal mixtures feature instead both entropic and energetic interactions, so that concentrations can be non-homogeneous at equilibrium even in the absence of chemical reactions. This is well described, for instance, by the Cahn-Hilliard theory of spinodal decomposition [14]. Recently, nonideal mixtures that undergo chemical reactions have attracted considerable attention due to their role in biology [15]. Reactions can affect the nature of phase separation and, when driven out of equilibrium, these

active systems exhibit rich phenomenologies. To be physicochemically justified and thermodynamically consistent, RD models need to express not only diffusion but also chemical dynamics in terms of nonideal chemical potentials [16,17]. Heuristic models of nonideal diffusion reactions have been considered, but use ideal chemical kinetics (mass action) and thus lack thermodynamic consistency [18–20]. Consistent models have been considered in Refs. [15,21–23]. However, they focus on unimolecular reactions, which cannot accommodate any spatial instability in the absence of energetic interactions. Extending these studies by considering multimolecular reactions is important because the instabilities that cause spatial organization can now arise from an interplay between chemical reactions and molecular interactions.

In this Letter, we consider thermodynamically consistent deterministic descriptions of generic nonideal mixtures of species undergoing diffusion and chemical reactions of arbitrary molecularity, driven out of equilibrium by external chemostats. Using linear stability analysis, we provide a rigorous classification of the possible instabilities and predict the conditions under which they arise. We find that they can be of two distinct types, which we call *E-type* and *R-type* instabilities. The former depends solely on the details of the intermolecular interactions, as in the Cahn-Hilliard theory of spinodal decomposition. The latter is controlled by the topology of the chemical reaction networks (CRNs), as in the Turing theory of instabilities in ideal mixtures. Unlike other classifications of RD instabilities [24,25] which focus solely on dynamics, ours is based on the underlying microscopic mechanism causing the instability. Importantly, we identify a wide class of CRNs where the instability can only be of E type. We also illustrate our findings with two specific models, each of them displaying one of two types of instabilities.

Chemical reactions and molecular interactions.—We consider an isothermal nonideal mixture at temperature T , composed of chemical species $\alpha \in \mathcal{S}$ which are reacting and diffusing within a solution of volume V . We partition the set of chemical species \mathcal{S} into two nonoverlapping subsets: the internal species $x \in \mathcal{X}$ and the chemostated species $y \in \mathcal{Y}$. The latter are exchanged with the external chemostats. Each chemical reaction $\rho \in \mathcal{R}$ is represented by the chemical equation



where Z_α is the chemical symbol of species $\alpha \in \mathcal{S}$, and $\nu_{+\rho}^\alpha$ [respectively (resp.) $\nu_{-\rho}^\alpha$] is the stoichiometric coefficient of species α in the forward (resp. backward) reaction $+\rho$ (resp. $-\rho$). We always use Einstein notation: repeated upper-lower indices implies the summation over all the allowed values for the indices. The set of internal species may include a nonreacting species ($\text{nr} \in \mathcal{X}$), defined by $\nu_{\pm\rho}^{\text{nr}} = 0$ for all $\pm\rho$. We assume that the chemostated species \mathcal{Y} are ideal and maintained at constant homogeneous concentrations, which result in the homogeneous chemical potentials μ_y . In practice, chemostats can drive chemical reactions far from equilibrium.

Turning to the dynamics, by combining dynamical density functional theory [26] and open CRN theory [27–29], the concentration fields of internal species $c_x(\mathbf{r}, t)$ evolve as

$$\partial_t c_x = D_x \nabla \cdot (c_x \nabla \mu_x) + S_x^\rho j_\rho, \quad (2)$$

with closure relations

$$\mu_x = \frac{1}{k_B T} \frac{\delta F}{\delta c_x}, \quad (3a)$$

$$j_\rho = j_{+\rho} - j_{-\rho}, \quad (3b)$$

$$j_{\pm\rho} = s_\rho e^{\mu_x \nu_{\pm\rho}^x + \mu_y \nu_{\pm\rho}^y}, \quad (3c)$$

where ∇ is the spatial gradient; k_B is the Boltzmann constant; μ_x (resp. D_x) is the nondimensional chemical potential (diffusion coefficient) of species x ; F is the Helmholtz free energy of the nonideal mixture; $S_x^\rho = \nu_x^{-\rho} - \nu_x^{+\rho}$ is the entry of the so-called stoichiometric matrix \mathbb{S} of the internal species (indexes x and ρ correspond to the rows and columns, respectively); j_ρ is the net current of reaction ρ expressed as the difference between the forward $j_{+\rho}$ and the backward $j_{-\rho}$ reaction flux; s_ρ is a positive preexponential factor that depends on the activation energy of the reaction ρ . The reaction fluxes $j_{\pm\rho}$ are defined in Eq. (3c) as the Arrhenius-like rates. We note that thermodynamically consistent currents could in principle allow for an additional dependence on the concentrations in $s_\rho(\mathbf{c})$, but this is rarely

considered and is thus omitted in our study. The diffusive contribution to the dynamics describes a pure gradient flow, and the only nonequilibrium drive stems from chemostats. In the absence of chemostats, the system relaxes to equilibrium. Dynamical models similar to Eqs. (2) and (3) have been recently considered in Refs. [21,22,30].

The Helmholtz free energy of the nonideal mixture reads as

$$F[\mathbf{c}] = k_B T \int d\mathbf{r} f(\mathbf{c}, \nabla \mathbf{c}) + F_{\text{chm}}, \quad (4)$$

where f is given in terms of the gradient expansion

$$f(\mathbf{c}, \nabla \mathbf{c}) = f_0(\mathbf{c}) + \frac{1}{2} K_{x,x'}(\mathbf{c}) (\nabla c^x) \cdot (\nabla c^{x'}), \quad (5)$$

the constant term F_{chm} is the contribution due to the ideal chemostated species, and $\mathbf{c} = (c_1, \dots, c_{|\mathcal{X}|})^\top$ (with $|\mathcal{X}|$ being the number of internal species). Here, f , f_0 , and $K_{x,x'} = K_{x',x}$ for $x, x' \in \mathcal{X}$ are model functions of the concentrations. Equation (5) is consistent with the free energy used in Refs. [31,32], and it can be straightforwardly extended to a more general form, including higher orders in gradients, as introduced by Cahn and Hilliard [14].

By using Eqs. (4) and (3a), the chemical potentials read as

$$\mu_x = \frac{\partial f_0}{\partial c_x} + \frac{1}{2} \frac{\partial K_{x',x''}}{\partial c_x} (\nabla c^{x'}) \cdot (\nabla c^{x''}) - \nabla \cdot (K_{x,x'} \nabla c^{x'}). \quad (6)$$

For ideal solutions, the free energy only comprises the entropic term $f = \sum_x c_x \ln c_x$, in which case one recovers a linear diffusion and mass-action kinetics in Eq. (2): $D_x \nabla \cdot (c_x \nabla \mu_x) = D_x \nabla^2 c_x$ and $j_{\pm\rho} \propto \prod_x c_x^{\nu_{\pm\rho}^x}$. For nonideal mixtures, the local free energy f_0 contains additional contributions, typically given as an expansion in powers of the concentrations [22], yielding nonlinear diffusion. If one considers homogeneous concentrations $\mathbf{c}^* = (c_1^*, \dots, c_{|\mathcal{X}|}^*)^\top$, we have $\nabla \mu_x^* = 0$ where $\mu_x^* = \partial f_0(\mathbf{c}^*) / \partial c_x$. Therefore, a homogeneous fixed point of Eq. (2) must satisfy the following steady-state condition:

$$S_x^\rho j_\rho(\{\mu_x^*\}, \{\mu_y\}) = 0. \quad (7)$$

Equation (7) shows that fixed points are determined by the chemical reaction contribution to the dynamics, which depends on (i) the stoichiometric coefficients, (ii) the chemical potentials of chemostated species μ_y , and (iii) the details of the local free-energy f_0 (i.e., including both entropic and energetic contributions). In contrast, for purely diffusive systems, each concentration is conserved, so that the homogeneous concentrations are fixed independently of the free-energy parameters.

Nature of instabilities: E type vs R type.—To analyze the stability of the homogeneous steady state, we consider small concentration perturbations around the homogeneous fixed point $c_x(\mathbf{r}, t) = c_x^* + \delta c_x(\mathbf{r}, t)$. Using the Fourier transform $\tilde{g}(\mathbf{q}) = \int d\mathbf{r} g(\mathbf{r}) \exp(i\mathbf{q} \cdot \mathbf{r})$, the perturbation of the chemical potentials in Eq. (6) can be written as

$$\delta\tilde{\boldsymbol{\mu}}(q) = \mathbb{M}(q) \cdot \delta\tilde{\mathbf{c}}(q), \quad (8a)$$

$$M_{xx'}(q) = \frac{\partial^2 f_0(\mathbf{c}^*)}{\partial c_x \partial c_{x'}} + q^2 K_{xx'}(\mathbf{c}^*), \quad (8b)$$

where $\delta\tilde{\boldsymbol{\mu}} = (\delta\tilde{\mu}_1, \dots, \delta\tilde{\mu}_{|\mathcal{X}|})^\top$ and $\delta\tilde{\mathbf{c}} = (\delta\tilde{c}_1, \dots, \delta\tilde{c}_{|\mathcal{X}|})^\top$.

Using Eq. (2), the evolution of a perturbation $\delta\tilde{\mathbf{c}}$ reads as

$$\partial_t \delta\tilde{\mathbf{c}}(q) = -q^2 \mathbb{A} \cdot \delta\tilde{\boldsymbol{\mu}}(q) + \mathbb{S} \cdot \delta\tilde{\mathbf{j}}(q), \quad (9a)$$

$$\mathbb{A} = \text{diag}(D_1 c_1^*, \dots, D_{|\mathcal{X}|} c_{|\mathcal{X}|}^*). \quad (9b)$$

By inserting $\mu_x = \mu_x^* + \delta\mu_x$ into Eq. (3c) and calculating the Fourier transform, we arrive at

$$\delta\tilde{j}_\rho(q) = s_\rho \left[\nu_{+\rho}^x e^{\mu_{x'}^* \nu_{+\rho}^{x'} + \mu_y \nu_{+\rho}^y} - \nu_{-\rho}^x e^{\mu_{x'}^* \nu_{-\rho}^{x'} + \mu_y \nu_{-\rho}^y} \right] \delta\tilde{\mu}_x(q). \quad (10)$$

Using Eq. (8b), we deduce that Eq. (9) becomes

$$\partial_t \delta\tilde{\mathbf{c}}(q) = \mathbb{B}(q) \cdot \mathbb{M}(q) \cdot \delta\tilde{\mathbf{c}}(q), \quad (11a)$$

$$\mathbb{B}(q) = -q^2 \mathbb{A} + \mathbb{C}. \quad (11b)$$

The elements of the square matrix \mathbb{C} are defined from Eq. (10):

$$C_x^{x'} = S_x^{\rho} S_{x'}^{\rho} \left[\nu_{+\rho}^{x'} e^{\mu_{x'}^* \nu_{+\rho}^{x'} + \mu_y \nu_{+\rho}^y} - \nu_{-\rho}^{x'} e^{\mu_{x'}^* \nu_{-\rho}^{x'} + \mu_y \nu_{-\rho}^y} \right], \quad (12)$$

where x and x' are the row and column index, respectively. We emphasize that the product structure of the Jacobian matrix $\mathbb{B}(q) \cdot \mathbb{M}(q)$ follows from the thermodynamically consistent description defining both diffusion and chemical fluxes in terms of chemical potentials.


Standard stability analysis [1] of Eq. (11a) implies that the homogeneous fixed point \mathbf{c}^* is unstable if at least one of the eigenvalues $\{\lambda_i\}$ of the Jacobian matrix $\mathbb{B}(q) \cdot \mathbb{M}(q)$ has a positive real part for a given wave number q . To avoid any divergence of the perturbations $\delta\tilde{\mathbf{c}}(q)$ at small wavelengths, we impose that all eigenvalues λ_i are negative as q tends to infinity [1]. In practice, this can be enforced by choosing appropriately $\{K_{xx'}\}$ in Eq. (5) which determines the cost of forming interfaces. This means that if we assume that $\text{Im}\lambda_i(q_0) = 0$, the condition for the homogeneous fixed point \mathbf{c}^* to become unstable, for at least one wave number $q_0 \neq 0$, can be expressed in terms of the determinant of the Jacobian matrix:

$$\det(\mathbb{B}(q_0) \cdot \mathbb{M}(q_0)) = (\det \mathbb{B}(q_0))(\det \mathbb{M}(q_0)) = 0. \quad (13)$$

Our analysis covers instabilities which typically induce stationary or transient patterns [24]. However, it does not cover instabilities often arising in homogeneous time oscillations and traveling waves [24], where simultaneously $\text{Re}\lambda_i(q_0) = 0$ and $\text{Im}\lambda_i(q_0) \neq 0$. In that case, the instability condition cannot be expressed in terms of the determinant of the matrix.

The condition in Eq. (13) shows that the instability can be caused by two distinct mechanisms: $\det \mathbb{M}(q) = 0$ or $\det \mathbb{B}(q) = 0$. Matrix \mathbb{M} depends on the free energy [Eq. (4)], and it also characterizes the purely diffusive system without reactions ($\mathbb{C} = 0$). The case $\det \mathbb{M}(q) = 0$ can only happen due to energetic interactions, since for ideal mixture \mathbb{M} is diagonal and positive. Thus, we refer to such an instability as an E type. In contrast, $\det \mathbb{B} = 0$ can happen in either ideal or nonideal solutions. The corresponding instability is not caused by energetic interactions, but instead by multimolecular chemical reactions. We refer to it as an R-type instability. Although matrix \mathbb{C} contains information on both the free energy (via the chemical potentials) and the stoichiometric matrix \mathbb{S} , the condition $\det \mathbb{B} = 0$ can only be met if \mathbb{S} satisfies certain conditions independent of the free energy, as we discuss below.

Restricted route to instability.—We now identify the specific class of CRNs where only instabilities of E type can arise. In this class, each reaction ρ interconverts m_ρ molecules of one specific internal species into m_ρ molecules of a different internal species, without constraints on the stoichiometry of the chemostatted species:



where $x \neq x'$, and there is no summation over x and x' (since they do not appear as repeated upper-lower indices). In Eq. (14), $\varepsilon_{x,\pm\rho}$ and $\varepsilon_{x',\pm\rho}$ can be either 0 or 1, so that every internal species is either a reactant or a product in a given reaction ρ . Furthermore, $m_\rho > 0$ is an integer number that can be different for each reaction ρ .

To prove that the CRNs (14) can only undergo E-type instabilities, we first demonstrate in Appendix A that the corresponding \mathbb{C} [Eq. (12)] has non-negative nondiagonal elements

$$C_{x,x'} \geq 0, \quad x \neq x' \quad (15)$$

and that the diagonal elements of \mathbb{C} satisfy the inequality

$$C_{x,x} \leq - \sum_{x' \neq x \in \mathcal{X}} C_{x',x}, \quad (16)$$

where the equality holds if and only if $\sum_{x \in \mathcal{X}} S_x^\rho = 0$, i.e., when the CRNs conserves the total concentration [27].

We then proceed to show that Eqs. (15) and (16) imply $\det \mathbb{B}(q) \neq 0$, which rules out R-type instability based on Eq. (13). To this end, we note that every eigenvalue of the matrix \mathbb{B} lies in the complex plane within (at least) one of a series of circles, referred to as Gershgorin circles [33]. In practice, each Gershgorin circle has a radius R_x defined as

$$R_x = \sum_{x' \neq x \in \mathcal{X}} |B_{x',x}| = \sum_{x' \neq x \in \mathcal{X}} C_{x',x}, \quad (17)$$

where we have used that \mathbb{A} is diagonal. Moreover, the center of each Gershgorin circle is located on the real axis (since \mathbb{B} has only real elements) at the point v_x given by

$$v_x = B_{x,x} \leq -q^2 D_x c_x^* - R_x, \quad (18)$$

where we used Eq. (16). As one can see from Eqs. (17) and (18), for $q > 0$ all Gershgorin circles are entirely located in the left complex plane. Thus, all eigenvalues of the matrix \mathbb{B} have a negative real part, so that $\det \mathbb{B}(q) \neq 0$. Combining this result with the condition in Eq. (13), it follows that the only way for the RD systems with reactions in Eq. (14) to entail any instability is $\det \mathbb{M}(q) = 0$, namely via a E-type instability.

Reactions described by Eq. (14) include pseudounimolecular ($\forall \rho: \nu_{\pm\rho}^y \geq 0$ and $m_\rho = 1$) and nonunimolecular reactions (for at least one $\rho: m_\rho > 1$). It is well known that ideal RD systems made of pseudounimolecular reactions cannot exhibit Turing patterns, as their dynamics is linear. In nonideal mixtures, energetic contributions to the free energy make the dynamics nonlinear even for pseudounimolecular reactions. Indeed, the reaction fluxes in Eq. (3c) and the matrix \mathbb{B} in Eq. (11b) explicitly depend on $\partial f(\mathbf{c}^*)/\partial c_x$ through the chemical potential μ_x [Eq. (6)], which could *a priori* trigger R-type instabilities. Yet, our result shows that nonlinearities stemming from molecular interactions can only create E-type instabilities for pseudounimolecular reactions. Importantly, this result extends to a special class of CRNs that also includes nonunimolecular reactions. We emphasize that, although such chemical reactions cannot generate R-type instabilities, their topology and rates strongly influence the location of the E-type instability, as they determine the homogeneous fixed points [Eq. (7)].

Illustrative examples.—We first consider the pseudounimolecular CRN in the inset of Fig. 1. In Appendix B, we derive its matrix \mathbb{C} and show that it satisfies Eqs. (15) and (16). Thus, this CRN belongs to the special class which admits only E-type instabilities. For the chemical potentials, we use the following expressions:

$$\mu_x = \mu_x^\theta + \log c_x + L_{x,x'} c^{x'} - K_{x,x'} \nabla^2 c^{x'}, \quad (19a)$$




FIG. 1. Inset: example of CRN that can only undergo E-type instabilities. Main: the blue curve is λ_+ for the RD system from the inset, the orange curve is λ_+^d for the corresponding purely diffusive system with same fixed points $\mathbf{c}^* = (0.12, 2.72, 0.10, 1.00)^T$. Parameters: $D_x = 1$, $\mu_1^\theta = \mu_3^\theta = \mu_{nr}^\theta = 0$, $\mu_2^\theta = -2$, $s_\rho = 10^{-3}$, $\chi = 1$, $k_1 = 0.5$, $k_2 = 0.1$, $\mu_{Y_1} = 1$, $\mu_{Y_2} = -1$ in arbitrary units.

$$\mathbb{L} = \begin{array}{c} X_1 \\ X_2 \\ X_3 \\ nr \end{array} \begin{array}{cccc} X_1 & X_2 & X_3 & nr \\ \left(\begin{array}{cccc} 0 & \chi & 0 & 0 \\ \chi & 0 & \chi & \chi \\ 0 & \chi & 0 & 0 \\ 0 & \chi & 0 & 0 \end{array} \right), & \mathbb{K} = \begin{array}{c} X_1 \\ X_2 \\ X_3 \\ nr \end{array} \begin{array}{cccc} X_1 & X_2 & X_3 & nr \\ \left(\begin{array}{cccc} k_1 & k_2 & 0 & 0 \\ k_2 & k_1 & k_2 & k_2 \\ 0 & k_2 & k_1 & 0 \\ 0 & k_2 & 0 & k_1 \end{array} \right). \end{array} \end{array} \quad (19b)$$

where μ_x^θ are the standard chemical potentials.

The matrices \mathbb{L} and \mathbb{K} describe the mean-field molecular interactions and the cost at forming interfaces, respectively. We numerically determine the homogeneous fixed points \mathbf{c}^* , from which we compute the matrices \mathbb{M} and \mathbb{B} according to Eqs. (8b) and (11b), respectively. In Fig. 1, for one of the fixed points we plot the eigenvalue $\lambda_+(q)$ of the matrix $\mathbb{B} \cdot \mathbb{M}$ which becomes positive over a finite range of q . We compared its behavior with the eigenvalue $\lambda_+^d(q)$ corresponding to pure diffusion system with $\mathbb{B}^d = -q^2 \mathbb{A}$ and $\mathbb{M}^d = \mathbb{M}$ at the same fixed point \mathbf{c}^* . Both $\lambda_+(q)$ and $\lambda_+^d(q)$ are positive in the same range of q , then vanish at the same point q_0 satisfying $\det \mathbb{M}(q_0) = 0$. This agrees with the scenario of E-type instabilities: it is sufficient to analyze the eigenvalues of the purely diffusive system to deduce the range of stability of the corresponding RD system. The model displays at least two additional homogeneous fixed points. One is stable $\lambda_+ < 0$. The other is unstable, and it is such that $\lambda_+(0) > 0$ and $\lambda_+(q)$ reaches a maximum at $q_m > 0$.

We now consider a nonideal version of the Brusselator model in the inset of Fig. 2. We use μ_1, μ_2 from Eq. (19a). This model does not satisfy the conditions in Eq. (14) and




FIG. 2. Inset (left): example of CRN undergoing a R-type instability. Main: the blue curve is λ_+ . Inset (right): the eigenvalues of the corresponding purely diffusive dynamics. Parameters: $D_1 = D_{nr} = 1$, $D_2 = 10$, $\mu_1^\theta = \mu_{nr}^\theta = 0$, $\mu_2^\theta = 9.2$, $s_\rho = 1$, $\chi = 0$, $k_1 = 0.1$, $k_2 = 0.05$, $\mu_{Y_1} = 9.9$, $\mu_{Y_2} = 1.1$, $\mu_{Y_3} = -27.6$, $\mu_{Y_4} = -18.4$ in arbitrary units.

thus can and does display an R-type instability. The right inset in Fig. 2 shows that while all eigenvalues of the purely diffusive process are negative, the eigenvalues of the RD Brusselator can be positive (Fig. 2), in stark contrast with the previous model. Indeed, since the Brusselator is not restricted to E-type instabilities, it can exhibit patterns in a regime where the corresponding purely diffusive system is stable.

Finally, we compare the results of numerical simulations for the two RD systems in Figs. 1 and 2. Figure 3 shows the steady-state 2D patterns of one of the chemical components. It is worth noting that the pattern in Fig. 3(a), for the RD system with reactions satisfying Eq. (14), is qualitatively analogous to the complete phase separation obtained




FIG. 3. The steady-state patterns for c_2 from numerical simulations in two-dimensional space with the periodic boundary conditions. (a) The unimolecular CRN with parameters from Fig. 1; (b) The Brusselator with parameters from Fig. 2. For simulations we used the py-pde package [34].

in purely diffusive dynamics. In contrast, the pattern of the Brusselator in Fig. 3(b) shows a striking different spatial organization which could not be reproduced by the corresponding purely diffusive system.

Discussion.—We characterized the nature of instabilities in thermodynamically consistent deterministic dynamics of nonideal RD systems in solution described by Arrhenius rates. We considered reversible reactions, but our results also hold for irreversible reactions. Extension to mixtures [15] is left for future work. Our decomposition in either E-type or R-type instability was shown for generic free energies and chemical reactions obeying the Arrhenius rate law Eq. (3c). It cannot be extended to rates laws where $s_\rho(c)$ depends on the concentration (due to the definition of the matrix \mathbb{B}). It can however be extended when $s_\rho(\mu)$ depends on the chemical potentials. Our proof that CRNs (14) can only undergo E-type instabilities however does not hold in this latter case and should be revisited. Future studies will be needed to study the thermodynamics (resp. fluctuations) of these systems, as recently done for ideal solutions [35–37] (resp. ideal [38] and nonideal RD systems [39]), and clarify their connection to heuristic active field theories including chemical degrees of freedom [40,41]. Turing instability in ideal solution requires different diffusion coefficients [12], but our condition for R-type instability does not. It would be interesting to explore if interactions could generate R instabilities inducing stationary patterns in models with the same diffusion coefficients. Our framework provides a tool to analyze how interactions promote or suppress the different types of instabilities, as recently done for models without thermodynamic consistency [42,43].

This research was funded by project ChemComplex (No. C21/MS/16356329) and project SMAC (No. 14389168) funded by Fonds National de la Recherche—FNR, Luxembourg, and by Project No. INTER/FNRS/20/15074473 funded by Fonds de la Recherche Scientifique—FNRS (F.R.S.—FNRS), Belgium and FNR (Luxembourg).

Appendix A: Proof of Eqs. (15) and (16) for CRNs that satisfy Eq. (14).—First, we recognize that the stoichiometric coefficients of the chemical equations (14) satisfy the following constraints:


$$\text{if } \nu_{x,\pm\rho} \neq 0 \text{ then } \nu_{x,\mp\rho} = 0, \quad (\text{A1a})$$

$$\text{either } \nu_{x,\pm\rho} = 0 \text{ or } \nu_{x,\pm\rho} = m_\rho, \quad (\text{A1b})$$

$$\text{either } \sum_x \nu_{x,\pm\rho} = 0 \text{ or } \sum_x \nu_{x,\pm\rho} = m_\rho. \quad (\text{A1c})$$

Second, we split the set of chemical reactions \mathcal{R} into two groups: internal and exchange reactions $\mathcal{R} = \mathcal{R}_{in} \cup \mathcal{R}_{ex}$. Internal reactions $\rho \in \mathcal{R}_{in}$ conserve the total concentration,

i.e., $\sum_x S_{x,\rho} = 0$ for $\rho \in \mathcal{R}_{\text{in}}$. The exchange reactions do not, i.e., $\sum_x S_{x,\rho} \neq 0$ for $\rho \in \mathcal{R}_{\text{ex}}$, and read as



namely, the internal species x is either a reactant or a product.

Third, we consider the nondiagonal elements $C_{x,x'}$ with $x' \neq x$. (i) Because of Eq. (A1c), $S_{x,\rho} = \pm m_\rho$ implies $\nu_{x',\mp\rho} = 0$. (ii) Because of Eq. (A2), $S_{x,\rho_{\text{ex}}} = \pm m_{\rho_{\text{ex}}}$ implies $\nu_{x',\rho_{\text{ex}}} = \nu_{x',-\rho_{\text{ex}}} = 0$. Hence, the terms of the summation over ρ in Eq. (12), namely,

$$\Xi_{x,x',\rho} = S_{x,\rho} s_\rho \left[\nu_{x',+\rho} e^{\mu_{x'}^* \nu_{+\rho}^{x'} + \mu_y \nu_{+\rho}^y} - \nu_{x',-\rho} e^{\mu_{x'}^* \nu_{-\rho}^{x'} + \mu_y \nu_{-\rho}^y} \right], \quad (\text{A3})$$

read as

$$\Xi_{x,x',\rho} = m_\rho s_\rho e^{\mu_{x'}^* \nu_{\pm\rho}^{x'} + \mu_y \nu_{\pm\rho}^y} \nu_{x',\pm\rho} \geq 0, \quad \text{if } S_{x,\rho} = \pm m_\rho, \quad (\text{A4a})$$

$$\Xi_{x,x',\rho} = 0, \quad \text{if } S_{x,\rho} = 0, \quad (\text{A4b})$$

$$\Xi_{x,x',\rho} = 0, \quad \text{if } \rho \in \mathcal{R}_{\text{ex}}, \quad (\text{A4c})$$

showing that all nondiagonal elements in Eq. (12) are non-negative, and thus proving Eq. (15).

Fourth, we consider the diagonal elements $C_{x,x}$. Because of Eqs. (A1a) and (A1b), $S_{x,\rho} = \pm m_\rho$ implies $\nu_{x,\mp\rho} = m_\rho$ and $\nu_{x,\pm\rho} = 0$. Hence, the terms of the summation over ρ in Eq. (12) become

$$\Xi_{x,x,\rho} = -m_\rho s_\rho e^{\mu_x^* \nu_{\mp\rho}^x} \nu_{x,\mp\rho} \leq 0, \quad \text{if } S_{x,\rho} = \pm m_\rho, \quad (\text{A5})$$

independently of whether $\rho \in \mathcal{R}_{\text{in}}$ or $\rho \in \mathcal{R}_{\text{ex}}$.

Fifth, we consider $\sum_x C_{x,x'}$. By using the splitting $\mathcal{R} = \mathcal{R}_{\text{in}} \cup \mathcal{R}_{\text{ex}}$, Eq. (A4c), and $\sum_x S_{x,\rho} = 0$ for $\rho \in \mathcal{R}_{\text{in}}$, we obtain

$$\sum_x C_{x,x'} = \sum_{\rho \in \mathcal{R}_{\text{ex}}} \Xi_{x',x',\rho} \leq 0, \quad (\text{A6})$$

proving Eq. (16).

Appendix B: Matrix \mathbb{C} for Fig. 1.—Here we derive the matrix \mathbb{C} for the example shown in Fig. 1. The coefficients $\nu_{\alpha,\pm\rho}$ can be written in matrix form:

$$\begin{aligned} \{\nu_{\alpha,+ \rho}\} &= \begin{matrix} & 1 & 2 & 3 & 4 & 5 \\ X_1 & \begin{pmatrix} 0 & 1 & 0 & 0 & 0 \end{pmatrix} \\ X_2 & \begin{pmatrix} 0 & 0 & 1 & 1 & 0 \end{pmatrix} \\ X_3 & \begin{pmatrix} 0 & 0 & 0 & 0 & 1 \end{pmatrix} \\ Y_1 & \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \end{pmatrix} \\ Y_2 & \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \end{pmatrix} \end{matrix}, \\ \{\nu_{\alpha,- \rho}\} &= \begin{matrix} & 1 & 2 & 3 & 4 & 5 \\ X_1 & \begin{pmatrix} 1 & 0 & 0 & 0 & 1 \end{pmatrix} \\ X_2 & \begin{pmatrix} 0 & 1 & 0 & 0 & 0 \end{pmatrix} \\ X_3 & \begin{pmatrix} 0 & 0 & 0 & 1 & 0 \end{pmatrix} \\ Y_1 & \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \end{pmatrix} \\ Y_2 & \begin{pmatrix} 0 & 0 & 1 & 0 & 0 \end{pmatrix} \end{matrix}, \end{aligned} \quad (\text{B1})$$

which gives us the stoichiometric matrix \mathbb{S}

$$\mathbb{S} = \begin{matrix} & 1 & 2 & 3 & 4 & 5 \\ X_1 & \begin{pmatrix} 1 & -1 & 0 & 0 & 1 \end{pmatrix} \\ X_2 & \begin{pmatrix} 0 & 1 & -1 & -1 & 0 \end{pmatrix} \\ X_3 & \begin{pmatrix} 0 & 0 & 0 & 1 & -1 \end{pmatrix} \\ \text{nr} & \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \end{pmatrix} \end{matrix}. \quad (\text{B2})$$

Using Eq. (12), (B1), and (B2), for Fig. 1 we find that

$$\mathbb{C} = \begin{matrix} & X_1 & & X_2 & & X_3 & & \text{nr} \\ X_1 & \begin{pmatrix} -s_1 e^{\mu_1^*} - s_2 e^{\mu_1^*} - s_5 e^{\mu_1^*} & & & & & & & 0 \end{pmatrix} \\ X_2 & \begin{pmatrix} s_3 e^{\mu_1^*} & -s_2 e^{\mu_2^*} - s_3 e^{\mu_2^*} - s_4 e^{\mu_2^*} & & & & & & 0 \end{pmatrix} \\ X_3 & \begin{pmatrix} s_5 e^{\mu_1^*} & s_4 e^{\mu_2^*} & & -s_4 e^{\mu_3^*} - s_5 e^{\mu_3^*} & & & & 0 \end{pmatrix} \\ \text{nr} & \begin{pmatrix} 0 & 0 & & 0 & & 0 & & 0 \end{pmatrix} \end{matrix}, \quad (\text{B3})$$

where the fixed point chemical potential μ_i^* for $i = 1, 2, 3$ corresponds to the species X_1, X_2, X_3 , respectively. From Eq. (B3) one can see that \mathbb{C} satisfies the properties in Eqs. (15) and (16). Notice that, as discussed in Appendix A, the exchange reactions—rates s_1 and s_3 —contribute only to the diagonal elements.

- *timur.aslyamov@uni.lu
 †francesco.avanzini@unipd.it
 ‡etienne.fodor@uni.lu
 §massimiliano.esposito@uni.lu
- [1] M. Cross and H. Greenside, *Pattern Formation and Dynamics in Nonequilibrium Systems* (Cambridge University Press, Cambridge, England, 2009).
- [2] L. A. Segel and J. L. Jackson, *J. Theor. Biol.* **37**, 545 (1972).
- [3] S. Kondo and T. Miura, *Science* **329**, 1616 (2010).
- [4] A. Mikhailov and G. Ertl, *The Frontiers Collection* (Springer, New York, 2017).
- [5] M. Z. Bazant, *Faraday Discuss.* **199**, 423 (2017).
- [6] A. M. Turing, *Phil. Trans. R. Soc. B* **237**, 37 (1952).
- [7] R. Lefever, *Phil. Trans. R. Soc. A* **376**, 20170365 (2018).
- [8] I. Prigogine and G. Nicolis, *Q. Rev. Biophys.* **4**, 107 (1971).
- [9] G. Nicolis and I. Prigogine, *Self-Organization in Nonequilibrium Systems: From Dissipative Structures to Order Through Fluctuations* (Wiley, New York, 1977).
- [10] L. Marcon, X. Diego, J. Sharpe, and P. Müller, *eLife* **5**, e14022 (2016).
- [11] X. Diego, L. Marcon, P. Müller, and J. Sharpe, *Phys. Rev. X* **8**, 021071 (2018).
- [12] P. A. Haas and R. E. Goldstein, *Phys. Rev. Lett.* **126**, 238101 (2021).
- [13] F. Brauns, J. Halatek, and E. Frey, *Phys. Rev. X* **10**, 041036 (2020).
- [14] J. W. Cahn and J. E. Hilliard, *J. Chem. Phys.* **28**, 258 (1958).
- [15] C. A. Weber, D. Zwicker, F. Jülicher, and C. F. Lee, *Rep. Prog. Phys.* **82**, 064601 (2019).
- [16] R. Lefever, D. Carati, and N. Hassani, *Phys. Rev. Lett.* **75**, 1674 (1995).
- [17] D. Carati and R. Lefever, *Phys. Rev. E* **56**, 3127 (1997).
- [18] S. C. Glotzer, E. A. Di Marzio, and M. Muthukumar, *Phys. Rev. Lett.* **74**, 2034 (1995).
- [19] M. E. Cates, D. Marenduzzo, I. Pagonabarraga, and J. Tailleur, *Proc. Natl. Acad. Sci. U.S.A.* **107**, 11715 (2010).
- [20] J. F. Lutsko and G. Nicolis, *Soft Matter* **12**, 93 (2016).
- [21] J. Kirschbaum and D. Zwicker, *J. R. Soc. Interface* **18**, 20210255 (2021).
- [22] D. Zwicker, *Curr. Opin. Colloid Interface Sci.* **61**, 101606 (2022).
- [23] J. Bauermann, C. A. Weber, and F. Jülicher, *Ann. Phys. (Berlin)* **534**, 2200132 (2022).
- [24] M. C. Cross and P. C. Hohenberg, *Rev. Mod. Phys.* **65**, 851 (1993).
- [25] T. Frohoff-Hülsmann and U. Thiele, arXiv:2301.05568 [Phys. Rev. Lett. (to be published)].
- [26] M. te Vrugt, H. Löwen, and R. Wittkowski, *Adv. Phys.* **69**, 121 (2020).
- [27] R. Rao and M. Esposito, *Phys. Rev. X* **6**, 041064 (2016).
- [28] F. Avanzini, E. Penocchio, G. Falasco, and M. Esposito, *J. Chem. Phys.* **154**, 094114 (2021).
- [29] F. Avanzini and M. Esposito, *J. Chem. Phys.* **156**, 014116 (2022).
- [30] A. M. Miangolarra and M. Castellana, *J. Stat. Phys.* **190**, 23 (2023).
- [31] S. Saha, J. Agudo-Canalejo, and R. Golestanian, *Phys. Rev. X* **10**, 041009 (2020).
- [32] E. Ilker and J.-F. Joanny, *Phys. Rev. Res.* **2**, 023200 (2020).
- [33] R. S. Varga, *Geršgorin and His Circles* (Springer Science & Business Media, New York, 2010), Vol. 36.
- [34] D. Zwicker, *J. Open Source Software* **5**, 2158 (2020).
- [35] G. Falasco, R. Rao, and M. Esposito, *Phys. Rev. Lett.* **121**, 108301 (2018).
- [36] F. Avanzini, G. Falasco, and M. Esposito, *J. Chem. Phys.* **151**, 234103 (2019).
- [37] F. Avanzini, G. Falasco, and M. Esposito, *Phys. Rev. E* **101**, 060102(R) (2020).
- [38] G. Falasco and M. Esposito, arXiv:2307.12406.
- [39] R. Tiani and U. C. Täuber, arXiv:2305.17726.
- [40] T. Markovich, E. Fodor, E. Tjhung, and M. E. Cates, *Phys. Rev. X* **11**, 021057 (2021).
- [41] E. Fodor, R. L. Jack, and M. E. Cates, *Annu. Rev. Condens. Matter Phys.* **13**, 215 (2022).
- [42] L. Menou, C. Luo, and D. Zwicker, *J. R. Soc. Interface* **20**, 20230244 (2023).
- [43] C. Luo and D. Zwicker, arXiv:2306.04882.